

Table V-8  
CERVIX UTERI CANCER (Invasive)

SEER INCIDENCE AND U.S. MORTALITY  
AGE-ADJUSTED RATES AND TRENDS<sup>a</sup>  
By Race/Ethnicity

SEER Incidence	Rate 2001-2005 <sup>b</sup>	Seer 13 Areas <sup>c</sup>	SEER 9 Areas <sup>d</sup>
	Rate per 100,000 persons	Trend 1996-2005	Trend 1996-2005
		AAPC (%)	AAPC (%)
RACE/ETHNICITY	Females	Females	Females
All Races	8.4	-2.8*	-3.8*
White	8.2	-2.4*	-2.6*
White Hispanic <sup>e</sup>	13.7	-3.6*	-
White Non-Hispanic <sup>e</sup>	7.0	-2.6*	-
Black	10.8	-3.9*	-3.8*
Asian/Pacific Islander	8.0	-4.8*	-
Amer Ind/Alaska Nat <sup>f</sup>	6.9	-2.3	-
Hispanic <sup>e</sup>	13.2	-3.5*	-
U.S. Mortality <sup>g</sup>			
	Rate 2001-2005	Trend 1996-2005 <sup>h</sup>	
	Rate per 100,000 persons	AAPC (%)	
RACE/ETHNICITY	Females	Females	
All Races	2.5	-3.1*	
White	2.3	-2.8*	
White Hispanic <sup>e</sup>	3.3	-3.2*	
White Non-Hispanic <sup>e</sup>	2.2	-2.7*	
Black	4.7	-4.6*	
Asian/Pacific Islander	2.2	-4.1*	
Amer Ind/Alaska Nat			
Total U.S.	2.7	-4.1*	
CHSDA Counties	3.7	-3.2*	
Non-CHSDA Counties	1.4	-	
Hispanic <sup>e</sup>	3.2	-3.1*	

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 3.3, April 2008, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

<sup>a</sup> Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Incidence data used in calculating the 2001-2005 rates are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

<sup>c</sup> The SEER 13 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia. The 1996-2005 AAPC trends are based on a Joinpoint analysis using diagnosis years 1992-2005.

<sup>d</sup> The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta. The 1996-2005 AAPC trends are based on a Joinpoint analysis using diagnosis years 1975-2005.

<sup>e</sup> Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. The 2001-2005 Hispanic and Non-Hispanic death rates exclude deaths from Minnesota, New Hampshire and North Dakota. The 1996-2005 Hispanic and Non-Hispanic mortality trends exclude deaths from Connecticut, Maine, Maryland, Minnesota, New Hampshire, New York, North Dakota, Oklahoma and Vermont.

<sup>f</sup> Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

<sup>g</sup> US Mortality Files, National Center for Health Statistics, CDC.

<sup>h</sup> The 1996-2005 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2005.

\* The APC is significantly different from zero (p<.05).